

Oculomotor Workshop

Timothy C. Hain, MD

Video Frenzel Goggles



Dr. Hain's system -- RealEyes Monocular

- Expensive (about \$2000)
- Good teaching tool
- Can do some things not easily done with optical system (i.e vibration test, hyperventilation, vertebral artery test, head prone test, cross-cover)

Optical Frenzel Goggles

Storz/German goggle



ICS goggle



- Inexpensive (about \$500)
- Portable – take to hospital
- A little limited – can't do vibration, head-forward or cross-cover. Some are dim.
- Can get hot, bulbs burn out and break

3

Frenzel variants that I don't like



- Any binocular goggle - -e.g. RealEyes xDVR, VOG-B100, VF-405 – heavy, fragile
- Any firewire goggle – good for computers, bad for bedside.
- Any “focus free” goggle - -e.g. InView – no focus means small and fuzzy.

4

Video Eye Movement Tests

- Saccades (slow, omn palsy, dysmetric)
- Spontaneous nystagmus
- Gaze testing
- Tests for unilateral loss
 - Vibration
 - Head-shaking
- Valsalva
- Hyperventilation


5

Slow saccades



6


Case



Source: Leigh/Zee

7


Case



Source: Leigh/Zee

8


Case



Source: Leigh/Zee

9

Case




Source of video: Preston Calvert, M.D.

10

INO (Internuclear ophthalmoplegia)


- Brainstem lesion of MLF
- Most commonly seen in MS
- Slowing of adducting saccades
- Overshoot of abducting eye
- A system that can visualize both eyes is best for INO, but too few INO's to justify trouble.



11

Oculomotor palsies

- Operators of ENG equipment may not think to do an oculomotor exam or check visual acuity. Useless output.
- Tests that displace viewing eye into paretic field eye may produce confusing results.




12

Case




13



Overshoot dysmetria


- Usually cerebellar lesion
- Occasionally parietic eye fixation
- Never peripheral vestibular lesion



15

Case

- 14 year old girl
- Very unstable gait
- headaches
- Darting eyes




16

Opsoclonus

- “dancing eyes-dancing feet” pediatric syndrome (Kinsbourne)
- Neuroblastoma
- Paraneoplastic syndrome
- West Nile

17

CASE



18

Case




19

Cases



20

Case



21

Spontaneous Nystagmus

- Acute vestibular disorders (V Neuritis, horizontal canal BPPV, Menieres, recent surgery) have strong horizontal “jerk” nystagmus.
- Normal people and chronic vestibular disorders have little or no nystagmus. Neural compensation for vestibular tone asymmetry is fast and effective.
- Most people can't “fake” nystagmus.
- Almost everything unusual is central.

22

Case – cross-cover test





23

Non-vestibular spontaneous nystagmus the common variants

Latent Nystagmus

- Found in persons with congenital esotropia
- changes direction according to viewing eye (Cross-cover test)
- Viewing eye beats laterally
- Intent to view controls direction (pseudoscope)
- Always have “lazy” eye

24




- ### Congenital Nystagmus
- One/1000 population
 - Present from early age
 - Usually worse in light
 - PT is not useful
 - Rehab significance is to avoid confusing it with central nystagmus or vestibular nystagmus.
- 27

- ### Non-vestibular spontaneous nystagmus: the common variants
- "Wrongly" directed primary position nystagmus
 - Downbeat
 - Upbeat
 - Torsional
- 28


Case

- Chiari (MRI)
- Cerebellar (especially remote effect) – get a CXR
- Idiopathic/drug

29



The cause



31

Case

- Smoking (slight)
- Paxil (slight)
- Wernickes
- BPPV variants ?
- Vestibular neuritis variants
- Central vertigo – Migraine ?



32

Direction ? Waveform




Something else was moving too



Oculo Palatal Myoclonus

- Fairly common disorder
- Pendular nystagmus
- Palatal myoclonus
- Triangle of Guillain Molleret



Gaze Testing

- Move finger to the limits of lateral gaze (bury sclera) – if can't bury, may have oculomotor palsy
- Move finger to limits of vertical gaze
- Do eyes reach end-gaze ?
- Is there end-gaze nystagmus ?
- Is there rebound nystagmus ?

36

Gaze Test: normal

- Minimal or no horizontal and upgaze nystagmus
- No down-gaze nystagmus in normal people
- No rebound nystagmus

37

Case (Cerebellar patient)

Patient BM
12/2003
Cerebellar
Degeneration

38

Rebound Nystagmus

- Nearly always cerebellar lesion
- Rarely congenital
- Method of separating out cerebellar GEN from sedative effect or congenital nystagmus

39

Vibration test



Vibration test



- Method: Apply 60-120 hz vibration to SCM, first one side, then the other. Shower massagers work well for this and are inexpensive. This is a Sunbeam/Oster shower massager
- Video Frenzel goggles – optical Frenzels don't work very well
- Compare nystagmus before and during

41

Vibration Induced Nystagmus

**Vibration Induced Nystagmus
in Unilateral Vestibular Loss**

(c) 2005 Timothy C. Hain, M.D.

42

Vibration Induced Nystagmus

- Unidirectional horizontal nystagmus strongly suggests contralateral vestibular lesion.
- Permanent nystagmus – never goes away
- Direction changing nystagmus is a normal variant.
- Vertical or torsional nystagmus is of uncertain meaning. Seems more common in BPPV.

Cherchi, M. and T. Hain (2010). Provocative Maneuvers for Vestibular Disorders. Vertigo and Imbalance: Clinical Neurophysiology of the Vestibular System. S. Eggers and D. S. Zee (Editors) . Elsevier.

43


Head-shaking test

- Method: 20 cycles of horizontal head rotation
- Frenzel goggles to monitor nystagmus prior to and following head-shaking.
- Positive – substantial change in nystagmus following head-shaking. Usually beats away from bad ear.



44

Head-shaking in person with left sided vestibulopathy



45

HSN – unilat comments

- SN, HSN and Vibration are all useful in detecting unilateral vestibular loss
- SN is seen acutely but vanishes over time.
- HSN is more sensitive to moderate loss than VN. However, it may appear and then vanish, or even go in wrong direction.
- Vibration is more dependable than HSN – never goes away.

Cherchi, M. and T. Hain (2010). Provocative Maneuvers for Vestibular Disorders. Vertigo and Imbalance: Clinical Neurophysiology of the Vestibular System. S. Eggers and D. S. Zee (Editors) . Elsevier.

46

Head Shaking

- Moderately useful test –
- About 75% localizing
- Absent in about 25%
- Small VOR nystagmus – good for bilaterals

Hain TC, Spindler J. Head-Shaking Nystagmus. in The Vestibulo-Ocular Reflex and Vertigo (Ed. Sharpe JA, Barber HO), Raven, 1993

47

Valsalva test

- Method: deep breath and strain
- Frenzel goggles to monitor nystagmus prior to and following HVT
- Positives suggest pressure sensitivity
 - Torsion - -SCD
 - Strong horizontal – horizontal canal fistula
 - Small amounts – not sure what this means



48



Hyperventilation test

- Method: 30 cycles of deep breathing
- Frenzel goggles to monitor nystagmus prior to and following HVT
- Positive – substantial change in nystagmus (other than DBN) following HVT.

50



Conclusion

Video Frenzel Goggles are the key to diagnosis of dizzy patients

- Oculomotor exam – far more sensitive with goggles
- Nystagmus → documents vertigo and localizes lesion
- Provocative testing → unilateral loss, SCD, VN nerve irritability

52

More details

Hain, T.C. Approach to the patient with Dizziness and Vertigo. Practical Neurology (Ed. Biller), 2002, 2007. Lippincott-Raven

More movies

www.dizziness-and-balance.com